

Name: Sonja Plasil
PI Name: Olivier George

Email: splasil@health.ucsd.edu
PI Email: olgeorge@health.ucsd.edu

Gene-Wide Association Study of Opioid-Induced Allodynia During Withdrawal in Heterogeneous Stock Rats

Sonja L. Plasil¹, Lani Tieu¹, Marsida Kallupi¹, Giordano de Guglielmo¹, Lieselot LG Carrette¹, Sierra Simpson¹, Jenni Kononoff², Adam Kimbrough¹, Lauren C Smith¹, Kokila Shankar^{1,2}, Alicia Avelar¹, Dana Conlisk², Molly Brennan¹, Sharona Sedighim¹, Brent Boomhower¹, Lisa Maturin¹, McKenzie J Fannon¹, Angelica Martinez¹, Caitlin Crook¹, Selen Dirik¹, Nathan Velarde¹, Paul Schweitzer¹, Selene Bonnet-Zahedi¹, Dyar N. Othman¹, Benjamin Sichel¹, Kwynn Guess¹, Beverly Peng⁴, Andrew S. Hu¹, Lucas E. Chun¹, Kristel Milan¹, Justin Lau¹, Yicen Zheng¹, Ashley Vang¹, Leah C. Solberg Woods³, Abraham A. Palmer^{1,4}, and Olivier George¹

¹Department of Psychiatry, University of California, San Diego, La Jolla CA 92093, USA

²Department of Neuroscience, The Scripps Research Institute, La Jolla, CA 92037, USA

³Department of Internal Medicine, Section on Molecular Medicine, Wake Forest University School of Medicine, Winston-Salem, NC 27157, USA

⁴Institute for Genomic Medicine, University of California, San Diego, La Jolla CA 92093, USA

Genetic variability accounts for 20-60% of the vulnerability to opioid use disorder (OUD) in humans. Pain is a driving force for opioid misuse and relapse. Preclinical rat studies suggest opioid self-administration (SA) could be independent from pain. This phenomenon produced challenges researching the link between OUD and pain. We hypothesize this discrepancy resulted from a lack of genetic diversity within animal models. Using heterogeneous stock rats (HS), a genetically diverse model relevant to OUD, we aimed to probe the relationship between opioid SA and hyperalgesia, perform gene-wide association studies of opioid-induced allodynia, and identify predictive gene variations. We hypothesize oxycodone SA and hyperalgesia are correlated, and this relationship strengthens or weakens based on the animal's addiction index (AI). A timecourse of mechanical sensitivity throughout withdrawal into protracted abstinence was completed using naïve HS and HS subjected to oxycodone SA. Von Frey behavior was performed at six timepoints following withdrawal: 4hrs, 12hrs, 24hrs, 1wk, 2wks, and 3wks. Results demonstrate oxycodone SA rats exhibit allodynia throughout protracted abstinence, and AI's correlated to the intensity and extent of mechanical hypersensitivity. Differentiation between naïve versus oxycodone animals, and high versus low AI animals was apparent at 12hrs and 24hrs, respectively. Low AI rats recovered to baseline by 1wk while high AI animals showed consistent hyperalgesia without baseline recovery. These data suggest oxycodone SA produces hyperalgesia following cessation, and mechanical sensitivity 12-24hrs into withdrawal may be predictive of the animals AI and oxycodone-seeking behavior. Assessment of gene variants predictive of oxycodone SA-induced allodynia is ongoing.